

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12Q 1/68	A1	(11) International Publication Number: WO 94/01582 (43) International Publication Date: 20 January 1994 (20.01.94)
(21) International Application Number: PCT/GB93/01452 (22) International Filing Date: 13 July 1993 (13.07.93) (30) Priority data: 9214873.3 13 July 1992 (13.07.92) GB (71) Applicant (for all designated States except US): MEDICAL RESEARCH COUNCIL [GB/GB]; 20 Park Crescent, London W1N 4AL (GB). (72) Inventor; and (75) Inventor/Applicant (for US only) : SIBSON, David, Ross [GB/GB]; 37 Grimsdells Lane, Amersham, Bucks HP6 HF (GB). (74) Agents: BIZLEY, Richard, Edward et al.; Hepworth, Lawrence, Bryer & Bizley, Gate House South, West Gate, Harlow, Essex CM20 1JN (GB).		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROCESS FOR CATEGORISING NUCLEOTIDE SEQUENCE POPULATIONS (57) Abstract A process for the categorization of nucleic acid sequences in which said sequences are linked to a population of adaptor molecules each exhibiting specificity for linking to a sequence including a predetermined nucleotide base, categorization of the resulting linked sequences being based upon selection for the base.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TC	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

PROCESS FOR CATEGORISING NUCLEOTIDE SEQUENCE POPULATIONS

This invention relates to a technique designed to facilitate study of the individual members of populations of nucleic acid sequences, particularly where the individual members of such populations may be present in widely varying amounts. This technique permits indexing of sequences.

Situations are increasingly arising in which it is necessary to study complex nucleic acid or polynucleotide populations. For example, it is now widely appreciated that an invaluable resource could be created if the entire sequence of the genomes of organisms such as man were determined and the information available. The magnitude of such a task should not, however, be underestimated. Thus, the human genome may contain as many as 100,000 genes (a very substantial proportion of which may be expressed in the human brain: Sutcliffe, Ann. Rev. Neurosci. 11:157-158 (1988)). Only a very small percentage of the stock of human genes has presently been explored, and this largely in a piecemeal and usually specifically targeted fashion.

There has been much public debate about the best means of approaching human genome sequencing. Brenner has argued (CIBA Foundation Symposium 149:6 (1990)) that efforts should be concentrated on cDNAs produced from reverse

transcribed mRNAs rather than on genomic DNA. This is primarily because most useful genetic information resides in the fraction of the genome which corresponds to mRNA, and this fraction is a very small part of the total (5% or less). Moreover, techniques for generating cDNAs are also well known. On the other hand, even supposing near perfect recovery of cDNAs corresponding to all expressed mRNAs, some potentially useful information will be lost by the cDNA approach, including sequences responsible for control and regulation of genes. Nonetheless, the cDNA approach at least substantially reduces the inherent inefficiencies resulting from analysis of repeated sequences and/or non-coding sequences in an approach which depends upon genomic DNA sequencing.

15

Recently, the results of a rapid method for identifying and characterising new cDNAs has been reported (Adams, M.D. et al., Science 252, 1991, pp 1651-1656). Essentially, a semi-automated sequence reader was used to produce a single read of sequence from one end of each of a number of cDNAs picked at random. It was shown, by comparing the nucleic acid sequences of the cDNAs (or the protein sequences produced by translating the nucleic acid sequence of the cDNAs) to each other and to known sequences in public databases, that each of the cDNAs picked at random, could be unambiguously classified. The cDNAs could be classified as being either entirely new or as corresponding, to a

20

25

greater or lesser extent, to a previously known sequence. cDNAs identified in this way were further characterised and found to be useful in a variety of standard applications, including physical mapping. Unfortunately, such a process is insufficient. The longer the process is pursued with any given population of cDNAs the less efficient it becomes and the lower the rate of identification of new clones. In essence, as the number of cDNAs which have already been picked rises, the probability of picking a particular cDNA more than once increases because of the wide range of abundancies at which different cDNAs are found, which abundancies can vary by several orders of magnitude. Thus, whereas some sequences are exceedingly rare, a single cDNA type may comprise as such as 10% of the population of cDNAs produced from a particular tissue (Lewin, B. Gene Expression, Vol. 2: Eucaryotic Chromosomes, 2nd ed., pp. 708-719. New York: Wiley, 1980). The need to avoid missing rarer species in any given population presents a considerable problem.

20

Various approaches have been tried in addressing the problem of increasing the efficiency of examination of a mixed nucleotide population, for example, such a population as is to be examined in human genome sequencing.

25

Thus, a standard PCR protocol can be used to amplify selectively cDNAs which are present at extremely low

levels, if there is information about the sequence of those cDNAs. If not, a primer specific to the desired cDNA cannot be constructed and the desired cDNA cannot be selectively amplified. The standard PCR method is therefore inadequate if it is desired to characterise a number of unknown genes.

Other approaches attempt to produce a more uniform abundance of the members of a population of cDNAs, so-called "normalisation methods". A first approach involves hybridization of cDNA to genomic DNA. At saturation, the cDNAs recovered from genomic/cDNA hybrids will be present in the same abundance as the genes encoding them. This will provide a much more homogenous population than the original cDNA library, but does not entirely solve the problem. In order to reach saturation in respect of the very rare sequences, it will be necessary to use huge quantities of cDNA, which need to be allowed to anneal to large amounts of genomic DNA over a considerable period of time. Furthermore, cDNAs which have homology to parts of the genome which are present in multiple copies will be over-represented.

A second approach exploits the second order reassociation kinetics of cDNA annealing to itself. After a long period of annealing, the cDNAs which remain single stranded will have nearly the same abundance, and can be recovered by

standard PCR (see Patanjali, S.R. et al., PNAS USA 88, 1991, pp. 1943-1947; Ko, M.S.H., NAR 19, no.18, 1991, pp 5705-5711). The methods disclosed in these two publications, however, suffer from notable disadvantages.

5 They are entirely dependent on the stringent physical separation of single stranded and double stranded DNA, require an elevated number of manual manipulations in each reaction, and necessitate protracted hybridisation times (up to 288 hours in the method of Patanjali et al.)

10

Yet a further approach in "normalising" a nucleotide population is described in copending British Patent Application No 91 15407.0, filed 17th July 1991 by MRC, and involves a PCR process in which a mixture comprising a
15 heterogenous DNA population and appropriate oligonucleotide primers is first formed and the DNA denatured, but before effecting a conventional PCR protocol the conditions are altered to allow the denatured strands of the more common DNA species to reanneal together, whilst avoiding annealing
20 of primers to the DNA strands. By this means, rarer species can subsequently be amplified in preference to the more common species.

25

This PCR normalisation method in general comprises the steps of:

(a) preparing a mixture comprising a heterogenous DNA

population and oligonucleotide primers suitable for use in a PCR process, in which the DNA is denatured;

(b) altering the conditions to allow the denatured strands of the more common DNA species to reanneal, while preventing the annealing of the primers to the DNA strands;

(c) further altering the conditions of the mixture in order to allow the primers to anneal to the remaining single-stranded DNA comprising the rarer DNA species; and

(d) carrying out an extension synthesis in the mixture produced in step (c).

Advantageously, the method consists of a cyclic application of the above four steps.

It will be appreciated that the conditions may be altered by the alteration of the temperature of the reaction mixture. However, any conditions which affect the hybridisation of complementary DNA strands to one another may be varied to achieve the required result.

Because the reannealing efficiency of any given DNA species will depend on the product of its concentration and time, the more abundant the sequence the greater the extent to which it will reanneal in any given time period. Once a

DNA species has reached a certain threshold concentration it will no longer be amplified exponentially, as a significant amount will have annealed to the double stranded form before the priming step. Thus, as each individual DNA species is amplified by the process to its threshold concentration, the rate of amplification of that species will start to tail off. Eventually, therefore, all DNA species will be present at the same concentration.

The length of the reannealing step will determine how much DNA is present at the threshold concentration. Preferably, therefore, the duration of the reannealing step will be determined empirically for each DNA population.

In the PCR normalisation process in general, the DNA primers may be adapted to prime selectively a sample of the total DNA population. By using primers which will only prime a sample of the population, only that sample will be amplified and normalised. The total quantity of DNA generated will thereby be reduced, which means that the cycling times can be kept low. This ensures that the method is applicable to complex DNA populations such as cDNA populations. In addition, a first primer can be used which is adapted selectively to prime a sample of the total cDNA population, and a second primer which is a general primer. Advantageously, the general primer is oligo dT (each primed cDNA will then be replicated in its entirety,

as the oligo dT primer will anneal to the poly-A tail at the end of the cDNA).

The present invention now provides a new process which
5 allows the study and identification of the individual
members of a mixed or heterogenous population of nucleic
acid sequences perhaps of varying abundance, for example,
the efficient identification of most cDNAs from a given
source tissue of a complex organism such as a human being.
10 The process of the invention is particularly adapted to
facilitate sorting (and investigation) of rarer sequences
in a population. Furthermore, nucleic acid sequences, for
example cDNAs as mentioned above, are produced in the
process in a way which makes them useful for new,
15 convenient and powerful approaches to a wide variety of
other applications.

Accordingly, the present invention provides a process for
categorizing uncharacterised nucleic acid by sorting said
20 nucleic acid into sequence-specific subsets, which process
comprises:

(a) optionally, initially subjecting said uncharacterized
nucleic acid to the action of a reagent, preferably an
25 endonuclease, which reagent cleaves said nucleic acid so as
produce smaller size cleavage products;

(b) reacting either the uncharacterized nucleic acid or, as the case may be, cleavage products from (a) with a population of adaptor molecules so as to generate adapted products, each of which adaptor molecules carries nucleic acid sequence end recognition means, and said population of adaptor molecules encompassing a range of such molecules having sequence end recognition means capable of linking to a predetermined subset of nucleic acid sequences; and

(c) selecting and separating only those adapted products resulting from (b) which include an adaptor of chosen nucleic acid sequence end recognition means.

In the above process, the adaptor molecules preferably comprise oligonucleotides in which single stranded ends of known nucleotide composition are present (the "nucleic acid sequence end recognition means" of the above process definition), said single stranded ends each exhibiting complementarity to a predetermined nucleic acid end sequence or end nucleotide so as to permit linkage therewith.

Of course, other forms of adaptor molecule can be envisaged. For example, adaptors can be chosen which are capable of specific reaction, preferably by covalent means, to any particular nucleotide or nucleic acid base or bases. Thus, advantage can, if desired, be taken of the existence

10

of unusual bases in certain nucleic acids, for instance, 5-hydroxymethylcytosine or thionucleotides. The nature of the linkage between adaptor and nucleic acid resulting from step (b) of the present process is irrelevant to success in the present invention, provided that such linkage is sufficiently strong to permit step (c) to be carried out, and further provided that the linkage is specific in a known way to only a known category of nucleic acid sequence ends or end nucleotides.

10

The process of the present invention can be applied to double stranded or single stranded nucleic acid materials, and there is no other particular limitation on the nature of the starting "uncharacterised" material which can be treated in the present process.

Preferably, step (b) of the present process is carried out with a population of adaptor molecules such that both ends of uncharacterised nucleic acid sequences or of the cleavage products from (a) can be adapted in a known way.

At least some of the adaptor molecules of the present invention can be structured so as to permit physical separation in step (c) of the present process by immobilizing adapted products on a solid phase. As an alternative, adaptor molecules can be used which comprise (in addition to their sequence end recognition means) a

known sequence permitting hybridization with a PCR primer. In such embodiments, the application of PCR techniques to a mixture of adapted products, using primers of preselected sequence, effectively enables one or more
5 predetermined subset(s) of sequences to be selected.

It will be appreciated that the starting nucleic acid material in the process of the present invention may already be in the form of a population of separate nucleic
10 acid sequences. Alternatively, even a lengthy continuous molecule, such as a complete chromosome, can be employed and, in order to produce a population of sequences for sorting and categorisation, the optional cleavage step of the present process, step (a), can be carried out.

15

If optional step (a) is effected, and if the reagent used for cleavage purposes is an endonuclease, this may be an enzyme specific to double stranded materials or it may be an enzyme which has the capability of cutting at a
20 recognized sequence on a single stranded product, depending upon the substrate uncharacterised nucleic acid.

Examples of suitable restriction endonucleases which recognise single stranded DNA and which also leave a
25 cleaved sequence overhang when cutting double stranded sequences (see later), which overhang is at least partly degenerate, include, BstNI, DdeI, HgaI, HinfI and MnlI.

Thus, for example, HgaI leaves a 5 base overhang starting 5 bases from the cut site, and MnlI cuts 7 bases away, but only leaving a 1 base overhang. Single stranded cleavage can also be achieved for enzymes which do not naturally cut single stranded DNA by annealing to the single strands an oligonucleotide containing a sequence for the recognition site for the enzyme, and which thus provides a partial double strand of sufficient length and nature for the enzyme to cut both strands.

One aspect of the power of the present pioneering process is the very generality of the materials which can be examined in applying, for the first time, an efficient technique for categorising and sorting nucleic acid sequences. Indeed, depending upon the number of stages in any sorting/selection process carried out (and the following description gives guidance as to various means whereby additional and further degrees of selection may be achieved), it is perfectly within the scope of preferred embodiments of the present invention to envisage adaptor molecules wherein the specificity is determined only by one nucleotide base available for linkage to an uncharacterised nucleic acid sequence end (eg specific to the final nucleotide).

As will be described hereinafter, the ability to sort/select to a high degree by sorting/selecting in

various stages is another great benefit of this invention. There is an exponential relationship between the number of such stages and the degree of sorting/selection.

5 One preferred aspect of the present invention is a process of categorising uncharacterised nucleic acid, which process comprises:

(a) subjecting said uncharacterised nucleic acid to the
10 action of a reagent, preferably an endonuclease which has cleavage and recognition site separated, which reagent cleaves said nucleic acid so as to produce double stranded cleavage products the individual strands of which overlap at cleaved ends to leave a single strand extending to a
15 known extent;

(b) ligating the cleavage products from (a) with adaptor molecules to generate adapted cleavage products, each of which adaptor molecules has a cleavage product end
20 recognition sequence and the thus-used adaptor molecules encompassing a range of adaptor molecules having recognition sequences complementary to a predetermined subset of the sequences of the cleavage-generated extending single strands; and

25

(c) selecting and separating only those adapted cleavage products resulting from (b) which carry an adaptor of known

recognition sequence.

The above preferred process of the present invention, aside from the general advantages of selection and sorting, additionally provides the extremely important advantage of being a means of indexing sequences because it enables a "marker" (in the form of a specific adaptor - see later) to be positioned at a predetermined site in any sequence. Furthermore, for the first time, sequence subsets can be produced in which not merely is something known about the individual sequence "ends", but also directionality in the sequences is established.

Generally, in the present invention it is convenient to use a population of adaptors simultaneously in step (b). Of course, if circumstances dictate, or if it is preferred for any reason, separate reactions may be performed with subsets of the total possible adaptor molecules required for "adapting" all possible sequence types.

In the present invention, "uncharacterised nucleic acid" is simply intended to mean any nucleic acid or population of nucleic acid sequences which is/are of partially or wholly unknown sequence.

As mentioned above, it is not significant to the generality of the present invention whether the uncharacterised

nucleic acid is double stranded or single stranded. However, in what follows a category of preferred embodiments of the invention is described in which Fok 1 is used as an endonuclease to generate nucleic acid fragments in accordance with step (a). If it is desired to use such an enzyme, or a similar enzyme, and the uncharacterised nucleic acid which it is desired to categorise consists of single stranded sequence material, such single stranded material can first be converted to double stranded sequences by methods known in the art (see, for example, Sanger, F. et al., Proc. Natl. Acad. Sci. 74, 1977, p5463-5467, Zoller, M.J. and Smith, M. Methods Enzymol. 100, 1983, p468-500, Gubler, U. and Hoffman, B.J. Gene 25, 1983 p263-269). The extent of strand overlap at the end of cleavage products resulting from step (a), in preferred embodiments, if carried out on double stranded material, may be as little as a single base, but is preferably two to ten bases, more preferably four to six bases. Preferably, as many as possible (at least 50%, and ideally 95% or more) of the cleavage products from (a) have each end overlapping in this way, and hence are capable of being "adaptored" by the preferred types of adaptor.

Preferred reagents which can be employed in step (a) are endonucleases, preferably Class II restriction endonucleases the cleavage sites with which are asymmetrically spaced across the two strands of a double

stranded substrate, and the specificity of which is not affected by the nature of the bases adjacent to a cleavage site.

5 When using the preferred oligonucleotide adaptor molecules of the invention, although any means of sequence-specific cleavage can preferably be used, most preferably the site of cleavage is not determined by sequence entirely at the ends of the fragments on the ends of different cleavage
10 products. Sequence specific chemical cleavage has been reported (Chu, B. C. F. and Orgel, L. E. Proc. Natl. Acad. Sci. p963-967 (1985)), but the preferred reagents are, as indicated above, a subset of Type II restriction endonucleases. This subset includes enzymes that have
15 multiple recognition sequences, enzymes that recognise interrupted palindromes and enzymes that recognise non-palindromic sequences. Type II restriction endonucleases of these types together cover a wide range of specificities, are readily available, and are highly
20 specific and efficient in their action (Review: Roberts, R. J. Nucl. Acids res. 18, 1990, p2331-2365). Some enzymes of the required type are listed in Table 1 below.

Table 1

	Enzymes with Multiple Recognition Sequences	Enzymes that Recognise Interrupted Palindromes	Enzymes that Recognize Nonpalindromic Sequences
5	Acc I	AlwN I	Alw I GGATC(4/5)
	All III	Bgl I	Bbs I GAAGAC(2/6)
	Aha II	BsaB I	Bbv I GCAGC(8/12)
	Ava I	BsaJ I	Bsa I GGTCTC(1/5)
	Ban I	BstE II	Bsm I GAATGC(1/-1)
10	Ban II	BstX I	BsmA I GTCTC(1/5)
	BsaA I	Bsu36 I	BspM I ACCTGC(4/8)
	Bsp1286 I	Dra III	Bsr I ACTGG(1/-1)
	BstY I	Drd I	Ear I CTCTTC(1/4)
	Clr10 I	EcoN I	Eco57 I CTGAAG(16/14)
15	Dsa I	EcoO109 I	Fok I GGATG(9/13)
	Eae I	Esp I	Gsu I CTGGAG(16/14)
	Gdl II	Nla IV	Nga I GACGC(5/10)
	Hae II	PflM I	Hph I GGTGA(8/7)
	HglA I	PpuM I	Mbo II GAAGA(8/7)
20	Hinc II	Sfi I	Mme I TCCRAC(20/18)
	NspB II	Tth111 I	Mnl I CCTC(7/6)
	NspH I	Xcm I	Ple I GAGTC(4/5)
	Sty I	Xmn I	Sap I GCTCTTC(1/4)
	Rsr II	Sfl I	SfaN I GCATC(5/9)
25			Taq II GACCGA(11/9)
			Tth111 ICAARCA(11/9)

30 Cleavage sites for enzymes which cleave outside of their recognition sequence are indicated in parentheses. For example, GGTCTC(1/5) indicates cleavage at:

35 5' ...GGTCTCN▼...3'
3' ...CCAGAGNNNNN▼...5'

40

45

Cleavages produced by some enzymes are blunt, while others produce a terminus with a single-stranded extension. Preferred enzymes fall into the latter category because these allow base specific ligation to be used for sorting
5 into subsets without having to produce single-stranded extensions from blunt ended termini.

As indicated earlier, a preferred endonuclease for use in the present invention is Fok 1.

10

An important feature of the present process is the use of adaptor molecules. The preferred adaptors generally have "overhanging" fragment recognition sequences which reflect or are complementary to the extending cleavage-derived
15 sequences which the adaptors are designed to react with. It is also preferred that such adaptors should end with a 5' hydroxyl group. The avoidance of a 5' phosphate group removes the risk of inappropriate ligation involving the adaptors. Alternatively, fragments to be adapted should
20 have their 5' phosphates removed and adaptors which have the potential to ligate to each other should be chosen so as to be separable by a means known in the art.

Adaptor molecules may also contain a portion permitting
25 specific sequence selection and separation (as in step (c) of the present process) when a sequence is attached to the adaptor. For example, an adaptor can carry biotin, thereby

permitting advantage to be taken of the biotin/avidin reaction in selecting and separating desired adapted molecules. Additionally, adaptors preferably comprise a known and selected sequence such that specifically adapted molecules can be amplified by known techniques (such as PCR) using a primer complementary to the core sequence.

Preferably adaptors in the invention are short double-stranded oligonucleotides which can be joined to the ends of cleavage products. They will have been chemically synthesised so that their sequence can be predetermined and so that large concentrations can be easily produced. They may also be chemically modified in a way which allows them to be easily purified during the process. As mentioned above, ideally their 5' ends will be unphosphorylated so that once joined to fragments the adapted end of the latter will no longer be able to participate in further ligation reactions.

[Preferably, all ligation reactions used in the invention will be catalysed by DNA ligase, since this enzyme is readily available and easy to use.]

It is preferred that the adaptor cleavage product end recognition sequences are on the 5' end of the longest oligonucleotide strand making up the preferred adaptor

molecules, are at least 3 nucleotides in length and with totally random bases at the single stranded position(s) two nucleotides in from the 5' end. This then allows selection to be performed both during the joining reaction and during
5 subsequent priming reactions. Then, because the final degree of selection is a result of the product of the degrees of selection achieved at these two stages, maximum selection can be achieved per adaptor/primer available (see later for further discussion).

10

Adaptor strand extensions on the 5' end of the longest oligonucleotide also facilitate the use of modified oligonucleotides for separation purposes. Preferably, the short oligonucleotide will be modified at its 5' end. This
15 has the double benefit of requiring just one modified oligonucleotide for all possible single-stranded extensions that are used, and also placing the modification at a position where it cannot interfere with ligation or subsequent priming reactions.

20

Although only one type of adaptor is required per ligation reaction, it is preferred that adaptors covering all possible reactions in a chosen subset of sequences be present, because then the opportunity for fragments in the
25 chosen subset to ligate to each other is minimised. It is also preferred that the chosen specific adaptor, carrying a predetermined recognition sequence, should not only be

different from the other adaptors in its single-stranded extension, but also different in the rest of its sequence since this allows orientation to be introduced which is useful in subsequent steps. It is therefore also preferred
5 that this adaptor has a modified oligonucleotide to facilitate its separation with the cleavage products to which it joins.

The process of the invention can be used to generate
10 categories or subsets of sequences by making some of the adaptors specific in some way, and selecting and separating as in step (c). In this way subsets of sequences can be provided depending upon the specific adaptor chosen, e.g. for use in subsequent nucleotide sequencing. This
15 facilitates, for example, the identification of a large population of sequences by permitting a rational approach to splitting such populations into subsets, each of which subsets can be examined in turn.

20 It will, of course, be appreciated that a subset generated in the present process can be regarded as a known fraction or specific proportion of all sequences in the original uncharacterised nucleic acid. A number of considerations can be used in determining the size and nature of any
25 particular subset. For example, subsets should not be too small because if the original nucleic acid provides a considerable number of different sequences, increasing the

- number of subsets also increases the number of adaptors and primers necessary and the number of experiments needed in order to categorise the nucleic acid. On the other hand, the person of skill in the art will need to tailor the subset size to give the required degree of resolution for any intended application. This is a matter of choice by the person of skill in the art, rather than general guidance.
- 10 Preferably there are only on average as many members of a subset as it would be convenient to identify in a later application of the present process, for example random picking for identification by sequencing. In this context, it is preferred that there be no more than a thousand members, preferably no more than five hundred members in a subset. A further point in relation to subset size is that it may be required to label simultaneously the subset for use as a probe. Preferably, in such a case, the total subset size should not exceed 500 kilobases, although this may be provided in a number of ways, e.g. 500 different 1 kilobase sized sequences or, for example, 1000 different 500 base sized sequences. Again, the matter is one of choice to the skilled operator.
- 25 It should also be remembered that adaptors are designed and required in the present process to ligate the fragments. In some circumstances, this can be driven kinetically e.g.

by the presence of large concentrations of adaptor. When this happens, subsets must be chosen so that the concentrations or diversity of fragments is such that the range of fragment ends available permits the desired adaptor concentrations to be attained.

This invention includes both the new adaptor molecules described herein, and also kits for performing a process of the invention which comprise a group of such adaptors designed to adaptor a predetermined group of nucleic acid sequences. In some embodiments, the present kits include a plurality of groups of different adaptors. The kits of the invention may also include, inter alia, a nucleic acid cleavage reagent, eg Fok 1 or other endonuclease as above, and/or PCR primers.

The new categorisation process of the invention will now be described further in relation to the accompanying drawings.

As will be appreciated, in the description that follows, including the specific Example, many individual features are referred to which are of broader applicability in carrying out embodiments of this invention than merely in performing any particularly described work.

In the drawings:-

Figure 1 is a schematic representation of a basic procedure behind the process of the invention for selection specific fragments - for simplicity it does not show all the adaptors which would be used in practice;

5

Figure 2 shows the cleavage behaviour of a preferred endonuclease for the invention, Fok 1;

10

Figure 3a illustrates the use of specific adaptors in accordance with the present process;

Figure 3b illustrates the selection of specifically adapted sequences after endonuclease digestion;

15

Figure 4 lists a range of adaptors and primers which can be used in the present invention; and

20

Figures 5a and 5b show specific predetermined priming (for amplification purposes) of specifically adapted molecules.

25

Figure 6 shows a gel obtained by agarose gel electrophoresis of samples categorised in the method of the invention as described in Example 2 hereinafter.

The process of the present invention is shown in outline in Figure 1. A nucleic acid or population of nucleic acids is

cleaved with a selected restriction endonuclease of known specificity and cleavage characteristics. The preferred example is Fok 1. The cleavage behaviour of Fok 1 is illustrated in Figure 2, and it will be seen that the single-stranded overhang or extension produced at the cleaved ends as a result of the action of Fok 1 can be any one of 256 possible four base sequences. However, at a given cleavage site the sequence will always be the same. Fok 1 is useful in this respect, both because it generates an overhang as illustrated in Figure 2 and also because its cleavage and recognition sites are separate.

The next stage in the process is the reaction of the population of nucleic acid sequence fragments resulting from endonuclease action with a population of adaptors. For the sake of simplicity, Figure 1 shows only the use of two specific adaptors, but it will be appreciated that for the purpose of categorising sequences in a mixed population or normalising such sequence populations, all possible adaptors specific for a predetermined subset of sequences must be used. This aspect will be referred to again hereinafter.

In Figure 1 one of the base specific adaptors is shown as being able to attach to a solid phase, and this attachment permits capture of fragments which have been adapted by the specific adaptor in question. Thereafter,

amplification techniques, e.g. PCR, may be used for base specific priming and amplification of the thus-isolated fragments.

5 When adaptoring is performed, it is important that all permutations of adaptor overhang be present or there will be the potential for hybrid molecules to form which are not representative of the original nucleic acid or nucleic acid population. Such hybrid molecules can form when fragments
10 which have not been specifically adapted ligate to each other or to the free ends of fragments which have been adapted in a specific way.

As can be seen most clearly in Figures 3a and 3b, it is at
15 the stage of adaptoring that the process of the invention introduces sorting or categorising of the fragments derived from endonuclease action. Obviously, any given adaptor will only react with a cleaved sequence where the overhang is complementary to the recognition sequence of the
20 adaptor. A degree of selection can thus be introduced at this point by preselecting one or more of the bases of the adaptor cleavage product end recognition sequence. Clearly, such specific adaptors are only capable of ligating with cleavage products or fragments in which the
25 overhang or extending sequence has a complementary base at the appropriate position or positions. This limits the proportion of the fragments resulting from endonuclease

activity which can be adapted by the specific adaptors.

A successful adapting stage in the present process depends upon the use of a restriction endonuclease which can produce any combination of bases in cleavage product ends. Clearly, if the overhanging bases were the same on all products or fragments resulting from cleavage, no selection would be possible.

As can be seen most clearly from Figure 3b, if a mixture of non-specific adaptors and specific adaptors is employed, and the specific adaptors are labelled or otherwise enabled in some way to be separated from the resulting mix, a sub-population of fragments is thereby automatically selected which has one or more predetermined bases in the overhang or single stranded extension at each cleavage site, which base or bases are predetermined by the choice of complementary base or bases in the recognition sequences of the adaptors.

20

As can be seen in Figure 3b, if specific adaptors are biotinylated, the adapted molecules which carry biotin residues can be bound to streptavidin-coated magnetic beads, washed and separated. Of course, other separation systems and labelling systems known in the art can be employed, and this is a matter of choice for the skilled reader (Uhlen, M., Nature 340, 1989, pp 733-744).

It will also be appreciated that although the technique illustrated in Figure 3b shows only one category of specific adaptor, which specific adaptor employs biotin/streptavidin in an art-understood way to isolate only those fragments to which the adaptor will bind, a number of specific adaptors can be employed simultaneously, provided such specific adaptors (and molecules to which they are attached when adaptoring has occurred) can be separated from each other.

In a subsequent stage of preferred process of the invention, if desired after the specifically adapted fragments have been separated, a further degree of selection can be achieved by copying or amplifying only selected subsets of the subset of specifically adapted nucleic acid fragments. Initial physical separation is, of course, not strictly necessary if a PCR-type process using only selected primers is employed. In any event, this further selection depends upon predetermined sequences in the core portion of the specific adaptors. Thus, once a particular set of specific adapted molecules has been isolated, for example by being bound to a solid phase using the biotin/streptavidin system, the nucleic acid fragments thus immobilised can be copied/amplified using a primer complementary to and specific for the core sequence of adaptors attached to immobilised nucleic acid fragments at

their ends remote from the sites of immobilisation. Such primers preferably extend by one or more specific extra bases into the adapted fragment.

5 As a result of this technique for further selection, only those adapted fragments can be copied which carry at their unbound ends the appropriate adaptor, preferably only those fragments having a sequence complementary to the chosen extra base or bases on the primer.

10

In this further selection technique it is highly preferred to use a polymerase which cannot significantly synthesise new strands from incorrectly annealed primers nor remove incorrectly annealed bases. Suitable enzymes include AMV reverse transcriptase (Kacian, D. L., Methods Virol. 6, 15 1977, p143), M-MLV reverse transcriptase (Roth, M. J. et al., J. Biol. Chem. 250, 1985, p9326-9335), DNA polymerase 1 Klenow (Exonuclease-Free) (Derbyshire, V., et al., Proc. Natl. Acad. Sci. U.S.A. 74, 1988, p5463-5467), genetically 20 engineered T7 DNA polymerase ((SequenaseTM) Tabor, S. and Richardson, C. C. J. Biol. Chem. 264, 1989, p6447-6458, U.S. Patent 4,795,699), and Taq DNA polymerase (Lawyer, F. et al., J. Biol. Chem. 264, 1989, p6427-6437).

25 Preferably, yet further selection can be achieved by selecting subsets of the resulting newly synthesised molecules. To do this, base specific priming can be

carried out as described above, except that the template for the primer in question is not the adaptor remote from immobilisation of the fragment, but rather the core portion sequence of the adaptor originally bound to the solid support. It will be appreciated that molecules obtained by copying at the first stage of selection subsequent to step (c) of the basic process of the invention are single stranded. A yet further stage of selection renders selected fragments double-stranded, and such fragments can be cloned by standard techniques see, for example, "Molecular Cloning", 2nd Edition Sambrook J., Fritsch, E. F., and Maniatis, T. CSH Press (1989).

The process of the invention can thus be used not merely to categorise molecules but also to select in a series of stages, hence "enriching" the amount available of any particular fragment. The final degree of enrichment depends upon the number of bases specifically predetermined at the adaptor and priming stages. By way of example, if two rounds of priming are employed, and if a specific adaptor is used which is specific for a single base and the first primer is specific for one base and the second primer is specific for two bases, the level of enrichment resulting from selection is 128 fold. This is because although each of the pre-determined or pre-selected bases which gives a single point of selection is one of four possibilities (giving a total number of permutations of 256), account

must be taken of the fact that each fragment has two ends which are capable of being adapted, thus dividing the degree of enrichment in half. It will be appreciated by those of skill that the same degree of enrichment can be achieved by using different combinations. For example, again assuming two rounds of priming, an adaptor which is specific for two bases only requires primers each specific for a single base in order to achieve 128 fold enrichment.

Not only is it important that the pre-determined bases should not be the same when determining or planning a particular degree of enrichment, it is also advantageous that the base permutations used for selection be distributed or "spread" in as many ways as possible since this minimises the number of primers and adaptors required. For example, a panel of 256 sequence subsets achieving 128 fold enrichment of a desired sequence each could employ a total of 256 base specific adaptors each of which is specific for one of the possible four bases in its "overhang". It is preferable, however, to use, for example, 16 adaptors which are specific for 2 bases each, together with 4 primers each specific for a single base at the 3' end of one of the adaptors and 4 more primers each specific for a single base at the 3' end of the other adaptor. This requires a total of only 24 adaptors and primers, corresponding to more than a ten fold reduction in the amount of these reagents required.

It will be appreciated that the shorter the individual members of any population of nucleic acid sequences of interest, the greater the percentage of nucleic acids which will only be cleaved once by the action of any chosen endonuclease. If most individual nucleic acid sequences in a population are not cleaved twice (or more) they will not be amenable to a full range of selection/enrichment choices in accordance with the principles set out above.

10 Schematically, the presence in the cleavage population of some fragments which do not lend themselves to adaptoring at both ends (as a result of a single cleavage only) is shown in Figure 3b. This problem can be addressed, if desired, by using two or more different restriction endonucleases, separately or together. Moreover, the present process is "adjustable" by the choice of an enzyme or enzymes where the spacing between recognition site and cleavage site is a favourable distance having regard to the size of the nucleic acid sequences in the chosen population

15 (if known).

20

Once subsets of nucleic acid sequences have been produced using the present process, they can be employed in a variety of ways. Labelled sequence subsets can be employed as probes, and libraries of subsets can be produced by cloning techniques. In addition, the subsets of sequences can themselves be probed after immobilisation using

25

standard techniques (see, for example, "Molecular Cloning" Maniatis et al, supra.)

The present process can be applied to a suitable source
5 material to produce subsets of restriction fragments, clones being picked at random for sequencing from the subsets, examining the subsets one by one. Once any particular subset becomes "exhausted" as an apparent source of new clones (or, more accurately, the rate of recovery of
10 new sequences drops) fresh subsets can be examined. A Poisson distribution can be used to describe the frequencies with which clones are picked at random from a subset in which members are present in equivalent proportions. Such a distribution is, of course, skewed
15 when members are not present in equivalent amounts, the skew being dependent upon the actual differences in the amounts. The observed distribution can be used to calculate the probability of picking at random new members of the subset, and this information can be used to decide
20 whether to persevere with a set for the purpose of picking new members. If the intention is to identify yet more members of a given subset, this may be more efficiently achieved by using a probe prepared using a pool of the already picked clones to identify clones which have not
25 already been picked and contain new sequences. The small sizes of the subset libraries which have to be probed in this way make this technique particularly convenient

compared to having to probe a library fully representative of the original starting nucleic acid.

It will be appreciated that the PCR normalisation technique described above can be applied to subsets produced by the process of the invention, rather than to the starting nucleic acid population or library, thus shifting the balance of nucleic acid sequences in individual subsets in favour of rarer sequences.

10

In investigating cDNAs from a tissue source, it is also useful to be able sequence such cDNAs from points other than their ends. In this way, a bias can be introduced in favour of potentially more interesting coding regions. The process of the present invention has the advantage of introducing such bias.

15

Fundamentally, however, the advantages of the process of the present invention include indexing sequences with consequent great advantages for understanding the structure of new sequences and mapping, and allowing cDNAs to be systematically picked from mRNAs prepared from an entire tissue or even an entire organism whilst increasing the "yield" of different sequences which can be obtained. This latter point is beneficial in recovering rarer sequences.

20

25

The invention will now be further described and illustrated

by means of the following Examples.

EXAMPLE 1

5 All oligonucleotides used in this Example were synthesised
Trityl on, using an ABI 380B DNA Synthesizer, according to
the manufacturers instructions. Purification was by
reverse phase HPLC (see, for example, Becker, C., R., et
al., J. Chromatography 326, p293-299 (1985)).

10

Human brain and adrenal tissues were obtained from a
mixture of 12 to 15 week menstrual age fetuses and then
snap frozen in liquid nitrogen before storing in bijou
bottles in a -80°C freezer. The two types of tissue were
15 used separately, directly from the freezer, to prepare cDNA
from which restriction fragments were generated for sorting
into subsets. 1g of each of the separate tissues were
homogenised, using an Ultra-Turrax T25 Disperser Janke and
Kunkel, IKA-Labortechnik, on ice in the presence of 4M
20 guanidinium isothiocyanate to solubilise macromolecules.
RNA was isolated from each homogenate by using
centrifugation to sediment it through cesium
trifluoroacetate. This was performed using the Pharmacia
kit according to the manufacturers instructions, except
25 that centrifugation was performed for 36 hours and the RNA
obtained was finally desalted and concentrated by
performing two ethanol precipitations in succession with

two 70% ethanol washes after each precipitation. In each case, polyA⁺ (mRNA) was isolated from 200 to 400 µg of the total RNA by binding it to magnetic oligodT coated beads (Dyna1). Solution containing unbound material was removed from the beads, which were washed, and then mRNA eluted directly for use. mRNA isolation was performed in accordance with the manufacturers instructions. Yields of RNA from the beads were between 1 and 3% of the total RNA. 2 to 4µg of the eluted RNA were used for cDNA synthesis. cDNA synthesis was performed according to the method of Gubler, U and Hoffman, B. J. Gene 25 p263 (1983) using a Pharmacia kit according to the manufacturers instructions. OligodT was used to prime the first strand cDNA synthesis reaction. The cDNA was purified by extracting twice with phenol/chloroform and then low molecular weight solutes including nucleic acids below ca. 300 bases were removed by passing the cDNA reaction mixture through a Pharmacia S400 spun column used according to the manufacturers instructions. Running buffer for the column comprised 10mM Tris-HCl, 1mM EDTA, 50 mM NaCl @ pH 7.5.

The column eluate was adjusted to 10mM Mg²⁺ and then the purified cDNA was restricted by the action of 1 unit per 10 µl of the endonuclease Fok I at 37°C for 1 hour, so that it would be able to accept adaptors.

The cDNA fragments were purified by two successive phenol/chloroform extractions followed by passing them through S400 spun columns as described above.

5 The adaptors used were oligonucleotides 5'
N₄N₄N₄N₄TCCTTCTCCTGCGACAGACA with the complementary strand
5' TGTCTGTCGAGGAGAAGGA and 5' AAN₄N₄TCTCGGACAGTGCTCCGAGAAC
or 5' TTN₄N₄TCTCGGACAGTGCTCCGAGAAC each with the
complementary 5' biotinylated strand
10 GTTCTCGGAGCACTGTCCGAGA. These were added to 25% of the
eluted material by incubating together 200 pmoles of the
mixture of double-stranded adaptors in the elution buffer
to which had been added MgCl₂ to 10mM, ATP to 10mM and 0.025
units/ μ l of T4 DNA ligase. The oligonucleotide 5'
15 biotinylated GTTCTCGGAGCACTGTCCGAGA, and whichever of the
complementary oligonucleotides with which it was used, each
comprised 1/32 of the molar proportion total adaptors. The
final reaction volume was 90 μ l which was heated to 65°C
for 3 minutes and then cooled to room temperature before
20 the ligase was added. Ligation was performed for 16 hours
at 12°C.

Two successive phenol/chloroform extractions were performed
to remove the ligase. The final aqueous phase was passed
25 through an S400 spun column (Pharmacia) as described above
except that the column was used with 10 mM Tris pH 8.3/50

mM NaCl.

The column eluate was adjusted to 25mM Mg²⁺, 0.5mM dNTPs in a final volume of 200 μ l. The mixture was placed in a thermocycler (Techne MW2) and heated to 78°C for 5 minutes. At this point 10 units of cloned Taq DNA polymerase (AmpliTag, Perkin Elmer) were added. This was followed by an incubation at 72°C for 10 minutes to fill in the unligated strand of the adaptor. After the second incubation 200 μ l of streptavidin coated magnetic beads (Dynal) prepared according to the manufacturers instructions were added to bind cDNA ligated to that of the oligonucleotides which was complementary to the 5' GTTCTCGGAGCACTGTCCGAGA biotinylated adaptor. Bead binding was allowed to proceed at 28°C for 30 minutes with mixing every 10 minutes.

Un-biotinylated cDNAs were washed from the beads with 400 μ l each of 2M NaCl twice, fresh 0.15 mM NaOH four times at 28°C for 5 minutes each, water twice and finally a buffer comprising 20 mM Tris pH 8.3, 50 mM NaCl, and 25mM Mg²⁺. The beads were then resuspended in 240 μ l of the final buffer including additionally 0.5 mM dNTPs and divided into 4x60 μ l.

25

Four of the 60 μ l aliquots, two from each tissue, were processed further specifically to prime and copy a subset

of the immobilised, adapted fragments. 2 pmoles of the primer 5' CTGTCTGTCGCAGGAGAAGGAA were added to each of two aliquots, one from each tissue. 2 pmoles of the primer 5' CTGTCTGTCGCAGGAGAAGGAG were added to each of the other two aliquots. 2.5 units of Taq DNA polymerase were added to each reaction and 16 cycles of alternate denaturation at 95°C for 30 seconds, annealing at 63°C for 2 minutes and polymerisation at 72°C for 3 minutes was performed to accumulate the selected single-strands in solution.

On completion of the DNA synthesis reactions a further 30 µl of resuspended beads were added to each reaction to remove the biotinylated fragments. The reaction was incubated at 28°C for 30 minutes mixing every 10 minutes to ensure that the biotinylated strands were bead bound. Each aqueous phase containing the newly synthesised strands was then removed and extracted with phenol/chloroform twice to remove the enzyme before being further purified by passing through an S400 spun column equilibrated with 10 mM Tris pH 8.3/50 mM NaCl as described above.

Rounds of PCR amplification of subsets of the selected fragments were performed by using the original primer in each case, together with one of the primers 5' GTTCTCGGAGCACTGTCCGAGAG or 5' GTTCTCGGAGCACTGTCCGAGAC. This simultaneously rendered the fragments double-stranded and increased the amounts of available material. It was

not known how many cycles of amplification would be required at this stage, since each primer pair would be expected to behave differently. It was therefore necessary directly to determine a suitable number empirically by using standard agarose gel electrophoresis to examine the reaction products after a given number of cycles. In some cases, to avoid the accumulation of non-specific products, it was necessary to perform an initial 5 cycles of amplification with both of the primers present at 2 pmoles each. All reactions were performed using 8 μ l or 12.5 % whichever was the larger but not exceeding 12 μ l of the column effluent above. Reaction conditions were adjusted to 20 mM Tris pH 8.3, 50 mM NaCl, 25mM Mg²⁺, 0.5mM dNTPs and 2.5 units of Taq DNA polymerase in a final volume of 40 μ l. Apart from when an initial amplification with 2 pmoles of each primer was performed, 20 pmoles of each primer were used. Cycles of amplification were performed at 95°C for 30 seconds, 65°C for 1 minute and 72°C for 3 minutes.

For the purposes of cloning, selected cDNA was amplified as described immediately above, except that the reaction was not monitored. Instead, the number of cycles which had previously been shown to just give rise to all observable products plus another 4 cycles were performed. In addition, an extra 72°C for 10 minutes incubation was performed after the last cycle.

The products of the reaction were then prepared for directional cloning. Water was added to adjust the final reaction volume to 60 μ l. Enzyme was removed by two successive phenol/chloroform extractions. The final aqueous mixture was passed through an S400 column as described above, except that it had been equilibrated with 10 mM Tris HCl pH 7.5, 50mM NaCl.

For directional cloning, advantage was taken of the different known sequences introduced at each end of the selected cDNAs by the adaptors in a modification of the method of Aslandis, C. and de Jong, P. J. (Nucl. Acids Res. 18, p6156 (1990)). Different cohesive ends were produced on each end by using the exonuclease activity of T4 DNA polymerase to resect from the 3' end, to the first T in each case. To 75 μ l or 75 % of the column eluate, whichever was least, were added 9.5 μ l of 100mM TrisHCl pH7.4, 100 mM MgCl₂, and 9.5 μ l of 0.5 mM dTTP. 16 units of T4 DNA polymerase were added and the reaction incubated in a water bath at 37°C for 30 minutes. The enzyme was removed by extracting with phenol/chloroform, twice successively. The salt of the final aqueous phase was adjusted by passing it through an S300 column (Pharmacia) equilibrated with 10 mM TrisHCl pH 7.4, 1 mM EDTA as described above.

The E.coli plasmid cloning vector pBluescript KS+ (Alting-

Meese, M. A. and Short J. M., Nucl. Acids Res. 17 p9494)
was prepared for accepting the resected cDNA by restriction
cleavage at the BamHI and HindIII sites and then adaptor
the resultant cohesive ends using the specific adaptors
5 produced by the oligonucleotide 5' AGCTCGGCTCGAGTCTG with
its partially complementary oligonucleotide 5'
GCGACAGACAGCAGACTCGAGCCG and the oligonucleotide 5'
GATCCGGCTCGAGT with its partially complementary
oligonucleotide 5' CCGAGAACACTCGAGCCG. Preparation of the
10 vector and adaptor were performed according to standard
procedures. Insertion of the cDNA was performed between
the BamHI and HindIII restriction sites. Recombinant
vectors were transformed into the host XL1-Blue (Bullock,
W. O. et al Biotechniques 5 p376-378 (1987)) by the method
15 of Hannahan, D. J. (Mol. Biol. 166 p577-580 (1983)).
Suitable standard controls for the ligations and
transformations were also included.

Post transformation procedures were as described in
20 "Molecular Cloning", 2nd Edition (Sambrook J., Fritsch, E.
F., and Maniatis, T. CSH Press (1989)). Colonies were
produced by plating onto X-gal/IPTG L-agar plates
containing 50µg/ml ampicillin and 10µg/ml tetracyclin.
Clear colonies were picked, each into a separate well of a
25 microtitre plate, containing 100µl of L-broth and 50µg/ml
ampicillin. Growth was allowed to occur for 16 hours at
37°C. 100µl of 50% or 30% glycerol was added to plates

which were archived at -20°C or -80°C, respectively.

Bacteria corresponding to those archived were used for preparing templates for sequencing by the dideoxy method (Sanger, F. Milklen, S. and Coulson, A. R. Proc. Natl. Acad. Sci. 74 p5463-5467 (1977)). Bacteria for this purpose were either grown on L-agar plates containing 50µg/ml of ampicillin, prepared at the same time as they had been grown in liquid culture, or after plating out from the archive. Alternatively, fresh liquid cultures were inoculated from the archive. In all cases, cDNA inserts were amplified for sequencing by PCR (Saiki, R. K. et al Science 239 p487-491 (1988)). PCR was either performed using bacteria directly added to the reaction, by a toothpick, or PCR was performed using 1/50th of the plasmid isolated by preparative methods (Holmes, D. S. and Quigley, M. Anal. Biochem. 114 p193 (1981)) from the bacteria in the liquid cultures or from the plates.

20 pmoles of each of the PCR primers 5' biotinylated GTAAAACGACGGCCAGT and 5' CGAGGTCGACGGTATCG were used in 40µl reactions containing 2.5mM Mg²⁺, 50 mM KCl, Tris-HCl pH 8.3 and 0.25 units of Amplitaq (Cetus). Reactions were performed at 95°C, for 1 minute, followed by 35 cycles at 95°C for 30 seconds, 60°C for 30 seconds and 72°C for 40 seconds. After the cycles, a final incubation at 72°C for 5 minutes was performed.

After PCR, standard agarose gel electrophoresis was used to determine which reactions had been successful. The biotinylated strands of successful reactions were then recovered for single-stranded sequencing by binding them to streptavidin coated beads (Dynal) and then washing, all according to the manufacturers instructions, except that the washing steps were either performed manually or performed automatically in the 96 well microtitre plate format using a Biomek robotic work-station attached to a side-arm loader (Beckman).

Dideoxy chain termination sequencing reactions were performed using the immobilised, biotinylated strands as templates and 2 pmoles of the oligonucleotide 5' CGAGGTCGACGGTATCG as primer. Reactions were performed using fluorescently-labelled terminators (Du Pont) or a fluorescein-labelled primer (Pharmacia) according to the manufacturers instructions. Reactions were analysed using automated DNA sequencers. A Genesis 2000 was used for the "Du Pont" reactions and an A.L.F. for the "Pharmacia" reactions. Bases were assigned for the Genesis 2000 reads using the manufacturers Base Caller software. Files of called bases were then transferred to a SUN Network from an Apple Macintosh computer which had been used for base calling. Raw data from the A.L.F. reads was directly transferred to a SUN network where bases were called using

the public domain "trace editor software" (TED). In both cases, files of called bases were entered into a SybaseTM database. Entering data entailed automatically removing vector and adaptor or linker sequences, but not editing ambiguous bases. After removal of the unwanted bases, files were automatically compared to other sequences in the cDNA database and the latest versions of the publically available databases, GENBANK and SWISSPROT. Searches were performed with the "basic local alignment search tool" (BLAST) (Karlin, S. and Altschul, S. F. Proc. Natl. Acad. Sci. 87 p2264-2268 (1990)).

It was expected that the amplification products of different subsets would appear qualitatively different because different cDNA fragments would be present in each. This was confirmed by agarose gel electrophoresis. Furthermore, amplification products were dependent on cDNA having been used. The amplification products were unlikely to be artefacts because their average size decreased on amplification. The spun columns used selected against material below 300 base pairs. This material would therefore have been expected to have been present in low amounts and therefore be the last to appear with amplification explaining the decrease in average size. The most usual amplification artefacts are multimers whose average size increases on amplification, and this was not observed.

Thirty five sequences, picked at random, were scored for the presence of a FokI site at the correct distance from the selected bases. 10 (28.5%) gave no data because of ambiguous bases. 15 (60% of the remainder) did not have such a site, while 10 (40% of the remainder) did have the expected FokI site. This is entirely as expected because Fok I can either be internal to a cleavage (in which case it appears in the selected fragment) or it can be external (in which case it is removed from a selected fragment). There is an equal likelihood of either possibility assuming perfectly random nucleic acid distribution. When different isolates of the same fragment in different clones were observed, these were always found to have the same sense with respect to the adaptors used and the vector. This is extremely unlikely by random assortment, and only likely as a result of the process of this invention. When observed sequences were found to correspond to already known sequences, the Fok I fragments selected were found to be as expected having regard to the bases used for selection, for example, the FokI fragment between bases 2642 and 3039 of DNA corresponding to the mRNA of the human amyloid A4 mRNA.

Example 2

25

Subsets of nucleic acid were prepared from cDNA as described in Example 1. cDNA was prepared from foetal

liver as described in Example 1, in addition to the foetal adrenal and foetal brain cDNA already described in that Example.

5 The restriction digestion, adaptoring and priming for the liver subset were performed as described in Example 1. In particular, the adaptor which was used for ligating to a specific set of fragments was 5'TTNNTCTCGGACAGTGCTCCGAGAAC and the primers used for priming of specific subsets during
10 PCR of the specifically adapted subsets were 5'CTGTCTGTGCGAGGAGAAGGAC (for the adaptor which had no ligation specificity for the required subset) and 5'GTTCTCGGAGCACTGTCCGAGAC (for the adaptor which was used to specifically select a subset of fragments). As a
15 control, subsets were identically prepared without the addition of cDNA. 20µl samples were taken after the final PCR reactions which produce the subsets and subjected to analysis by agarose gel electrophoresis, the nucleic acid being detected by ethidium bromide. The results are seen
20 in Figure 6.

In Figure 6, the significance of the lanes is as follows:

	Lane 1	-	marker
25	Lane 2	-	as 6, no cDNA added
	Lane 3	-	as 7, no cDNA added
	Lane 4	-	as 8, no cDNA added

48

- | | | | |
|----|--------|---|--|
| | Lane 5 | - | size marker : 2176, 1766, 1230, 1033, 653, 517, 453, 394, 298, 234 base pairs respectively, uppermost to lowermost visible bands |
| 5 | Lane 6 | - | fetal brain cDNA subset, adaptor specific for tt, specific adaptor primer specific for g, non specific adaptor primer specific for c |
| 10 | Lane 7 | - | fetal adrenal cDNA subset, adaptor specific for aa, specific adaptor primer specific for g, non specific adaptor primer specific for g |
| 15 | Lane 8 | - | fetal liver cDNA subset, adaptor specific for tt, specific adaptor primer specific for g, non specific adaptor primer for g |

It can be seen from lanes 2 to 4 and 6 to 8 of Figure 6 that unless cDNA was added no amplification was observed, illustrating that de novo amplification, for example by primer dimerisation, is not a problem. Furthermore, in addition to the other products, a strong specific band was observed in the foetal liver and a different strong specific band in the foetal adrenal samples. Several weaker bands were observed in the foetal brain sample. This provided an opportunity to test the specificity of the sorting procedure.

The unused material was cloned as described in Example 1, and then individual clones were sequenced as described in Example 1. The sequence files from each of the subsets were separately copied from the database. The sequences from a given subset were then compared to each other using icaass, a global similarity comparison algorithm from the software icatools (Parsons, J et al CABIOS 4 p367-271 (1988). This determined how often the same restriction fragment from the sorted cDNA was observed. By comparison to the BLAST searches, it could be deduced that assuming no significant cloning bias, 21.4% of the liver subset was the Fok 1 restriction fragment at the end corresponding to the carboxyl terminus of serum albumin (bases <2080 on Genbank accession no = L00132), and that this was the major band observed in the agarose gel. The size of the band is consistent with this deduction. There are 6 possible Fok 1 restriction fragments from serum albumin (which is an abundant transcript) so substantial selection by the sorting procedure had occurred. Similarly, the abundant restriction fragment in the adrenal subset (6.4%) originated from the mitochondria within the region of the gene for NADH-ubiquinone oxidoreductase chain 4 (bases >11711 on Genbank accession no = J01415). No restriction fragments dominated the brain subset. These results are also consistent with the agarose gel analysis and again indicate that selection has occurred.

Cloning was performed with an orientation with respect to the selective and the non-selective adaptors. This provides an additional test of the sorting procedure because, if no selection was occurring, when a fragment was observed more than once its orientation would be expected to be random, with 50% of instances in one orientation and 50% in the opposite orientation. In fact, the results given in Table 2 below show that when a fragment occurs more than once, as determined by the icaass comparison, in none out of 40 cases did antisense matches occur. This is highly improbable unless selection has introduced an orientation which has been maintained during the directional cloning. Selection was achieved through adaptor ligation and through priming, thus each of these must have had a high degree of specificity during the selection. The specificity of the ligation is remarkable given that the oligonucleotides in the ligation reaction covered every possible combination of bases at each of the four possible base positions closest to the 5' end of the non-specific adaptor. As a comparison the sequencing and analysis was performed as described in Example 1, except that conventional, commercial cDNA libraries which had not been sorted or directionally cloned were used as the source of the cDNA clones, and appropriate sequencing primers were used. The results for these comparisons are also shown in Table 2. It can be seen that antisense matches are commonly observed at a frequency of 10.5% and 36.3% in the

51

adult liver (Clontech Laboratories, Inc Cat No HL1001b) and
adult brain cortex (Clontech Laboratories, Inc Cat No HL
10036) libraries, respectively.

5

10

15

20

25

Table 2

Comparison of frequency classes and the proportion of antisense matches between cDNA subsets and total cDNA libraries.

Tissue	Subset Brain	Subset Brain	Subset Adrenal	Subset Adrenal	Subset Adrenal	Subset Adrenal	Subset Adrenal	Total cDNA Libraries
Adaptor specificity	aa	aa	aa	aa	aa	aa	tt	Adult Liver
Primer for specific adaptor	a	g	a	a	g	g	a	Adult Brain
Primer for non specific adaptor	g	g	g	g	g	g	g	(cortex)
Frequency class								
1	183	117	58	163	38	226	341	
2	14	2	2	2		15	19	
3	3	2		2	1	4	3	
4							1	
5								52
6								
7								
8								
9								
10								
11								
12								
13								
14								
15								
16								
% unique patterns	90.91%	95.28%	96.77%	90.27%	95.12%	89.42%	84.62%	
Antisense matches	0	0	0	0	0	10.5%	36.3%	

Using the present process, once restriction fragments from an original nucleic acid have been sorted into subsets they can be advantageously manipulated and analysed by the full repertoire of molecular genetic or molecular biological techniques. They are also useful for a wide variety of applications which take particular advantage of their sorted nature. Compared to the situation in the original population, some of the useful types of property conferred on the fragments by sorting are that their relative abundance is higher in the sorted population, that the sequences at the ends of all of the members of a sorted population are the same and therefore indexed, that the members of a sorted population form a unique and discrete set, that the subset produced from one type of nucleic acid will be comparable to the same subset produced from a related nucleic acid, and that by separately recognising the two ends of the fragments in a subset they acquire known directionality.

20

The application of the method for sorting restriction fragments produced from cDNAs in the human genome sequencing project is amongst the most technically demanding examples of the application of the present process which can be envisaged. Single-stranded RNA first had to be converted into double-stranded cDNA. Furthermore, the fragments produced from cDNA were present

25

in a wide variety of different abundances, reflecting the original composition of the mRNA population. Nevertheless, sorting was successfully achieved, thus illustrating the robustness of the method.

5

The known directionality of sorted fragments makes it possible, in principle, to use fragments directly for probing by PCR. This would require that the same fragment be isolated from two identical subsets, except that the
10 core sequences of the adaptors used for each subset would be different. The fragment from one set would then be immobilised at the 5' end of one of its adapter sequences and then annealed to the target. No annealing between target and fragment could occur at sequences which
15 corresponded to the adaptors. A DNA polymerase, with a 3' exonuclease as its sole nuclease activity, would then be used to resect the unannealed 3' ends until the first complementary annealed base was reached, at which point the reverse complement of the non-immobilised adaptor sequence
20 would become copied into the target. After resecting, the nucleic acids would then be heat denatured, simultaneously inactivating the enzyme, and the immobilised fragments removed to leave target which had incorporated the non-immobilised adaptor sequence. Copies of the target could
25 then be made from a primer which corresponded to the incorporated adaptor. Target would then be denatured and annealed to the complementary strand of the fragment from

55

the second subset. This fragment would also have been immobilised at its 5' end. Resecting would be performed as before. This time, the reverse complement of adaptor sequences on the target would become incorporated onto the immobilised fragment. Fragments which owed their origin specifically to the target could then be exclusively amplified by PCR using a primer which corresponded to the current immobilised 5' end and a primer which corresponded to the previous immobilised 5' end. It may be possible to use more than one type of fragment at once in this approach. Furthermore, exact complementarity between probe and target may not be required, thus allowing polymorphisms to be detected.

The fact that members of sorted populations form unique and discrete sets could be used more conveniently to identify fragments of interest. Fragments would be sorted into subsets such that any member of a subset could be detected using a suitable hybridization probe. Probing the subsets would identify in which subset a fragment of interest could be found. The fragments in that subset would then be probed to find the one of interest. This two step approach would be easier than probing all possible fragments at once.

25

The principle that members of sorted populations form unique and discrete sets can also be exploited during

sequencing projects. Two or more different restriction endonucleases could be used to produce two or more corresponding sets of subsets in the knowledge that overlapping fragments must be produced between the sets of subsets. The members of each subset would then be sequenced. The degree of sorting would have been determined in advance to produce a number of fragments per subset that would allow each of the members to be easily identified. This would minimise the need repeatedly to read the same sequences, as commonly occurs during other approaches to sequencing projects. In addition, fewer gaps would be expected in the final overlapping sequence. It is often desirable to read a given sequence several times so that bases corresponding to ambiguities can be confirmed. This could easily be achieved by reading different isolates of the same fragment. Gaps would occur in the sequence either because inserts were too long to be sequenced or because fragments were of an inappropriate size to be successfully sorted. It would be possible conveniently to fill such gaps by using the known flanking sequence to predict in which subset a fragment produced by a different restriction endonuclease could be found for the purpose of extending the sequence. Fragments from the vector would also be present, but they could easily be avoided since it could be predicted in which subset they would occur and they could be identified from their size.

The fact that fragments are in a higher relative abundance in a sorted subset than in the original population is likely to benefit applications which use nucleic acid hybridization probes. In such applications, sensitivity is
5 determined by "signal to noise" ratios. This becomes critical in applications where not all of the probe is specific for a particular target and can therefore contribute to background (thus reducing sensitivity).

10 A limit is therefore set on the size of a nucleic acid or the number of different nucleic acids which can be used simultaneously as a probe, which in turn reduces the possible range of applications. These limits can be overcome by using, separately as probes, sorted subsets of
15 the original intended probe. In this way, probing applications are made possible that otherwise would require detailed subcloning and/or characterisation of the original nucleic acid. Examples of nucleic acid types known in the art which could be used as probes or targets in any
20 possible combination in this way include:- cDNA fragments, clones or libraries; lambda genomic clones or libraries; cosmid genomic clones or libraries; P1 genomic clones or libraries; YAC genomic clones or libraries; sorted chromosomes or products of sorted chromosomes; chromosome
25 specific libraries; products or libraries of material from microdissected chromosomes; products of co-incidence cloning or libraries of co-incidence clones; products of IRS-PCR or

libraries of these products.

In general, the aim would be to detect sequences in common between the possible pairings. It would be particularly
5 helpful when there were sequences of widely different abundance in the probe and/or target, since it would allow simultaneous detection of sequences that would not normally be possible as a result of wide differences in their signal strength. A classic example of such a
10 situation is when differential screening of cDNA libraries is being performed. Typically, clones in two different cDNA libraries are detected using a probe prepared from the cDNAs used to make one of the libraries. The strength of the signal from a given clone indicates its abundance in
15 the library to which the probe corresponds. Many clones will be an abundance which is too low to be detected. Appropriate enrichment of the probe in subsets through sorting would allow all clones to produce a measurable signal.

20

Given that, by sorting, the enrichment of fragments from a probe can be tailored so that even the least abundant sequence in a target can be detected, that the members of a sorted population form a unique and discrete set, and
25 that the subset produced from one type of nucleic acid will be comparable to the same subset produced from a related nucleic acid, it is possible to fingerprint individual

members of libraries of interest for the purpose of clone identification or mapping. This also requires that subsets are discretely, partially overlapping, which, as a result of both ends of any given fragment being independently
5 sorted, they are. Each subset from a probe would be used in turn to detect clones in one or more libraries of interest. Each clone would then acquire a signature determined by the subsets which detected it. Resolution would depend on how many features were in a subset, how
10 many subsets were produced and by the number of clones for which a signature was determined. A sufficient number of subsets could be produced by preparing them from more than one enzyme so that each clone could be given a unique signature through which it could be identified. Clones
15 with the same signature could be assumed to have the same insert, while for the purposes of physical mapping, overlapping signatures would signify overlapping inserts.

A common requirement during genetic mapping projects, for
20 example in clinical genetics, is to screen for bases which are polymorphic between individuals, so that the fate of the genomic region in which the polymorphism are found can be monitored through generations for linkage analysis. Restriction site polymorphisms are useful in this respect.
25 Base polymorphisms can result in individuals differing with regard to whether they possess a detectable restriction site. The twin principles that the members of a sorted

population form a unique and discrete set and that the subset produced from one type of nucleic acid will be comparable to the same subset produced from a related nucleic acid provide a convenient means of detecting clones which themselves can be used as probes to detect polymorphisms at restriction sites between individuals. Clones would be prepared from the fragments in a subset from one individual. These would then be probed using the fragments which were used to produce the clones, and also separately probed using fragments from the corresponding subset prepared from a different individual. Clones which could detect restriction site polymorphisms between the individuals would be identified as those which failed to give rise to a signal when the subset from the individual from which they did not originate was used as a probe as compared to when the subset from the individual from which they did originate was used as a probe. The polymorphism would be in a site for the enzyme which was originally used to prepare the subset. Clones of genomic or cDNA could be used for the application. Use of cDNA clones would have the advantage that it would be known that the restriction site polymorphisms were being detected in actual genes, whose fate could then be followed through generations.

It would be particularly advantageous if the individuals between whom the polymorphisms were detected as described above were those who corresponded to the grandparents in a

back-cross. 50% of the individuals in the F2 generation could then be scored for the polymorphisms detected, assuming that back-crossing had been performed equally to each of the original grandparents. It would also be useful
5 if the nucleic acid from each of the F2 individuals was separately sorted and amplified. Polymorphisms could then be scored in the amplified F2 subsets corresponding to these in which the polymorphisms were detected between the original grandparents. This would have the triple
10 advantage that the usually finite back-cross resource would have been immortalised for this purpose, that the present or absent nature of signal from each sample would allow many samples to be conveniently probed simultaneously after spotting them onto a gridded array, and that the target,
15 where present, would have been enriched in a subset so that it could be more easily detected. This could be of particular value in plant or animal breeding.

In view of the above, it will be understood that the
20 invention includes any use of a method of the invention or of a population of sequences, sorted or categorised in accordance with the invention, where said use takes advantage of any one or more of:

25 (a) the relatively higher abundance of sequences in a sorted population;

(b) the "indexing" of sorted sequences by the use of adaptors;

(c) the fact that members of a sorted population form a
5 unique and discrete set;

(d) the fact that a given subset produced from one nucleic
acid source material will be comparable to the
correspondingly "adaptored" subset from a related nucleic
10 acid; and,

(e) as a result of adaptoring sequences acquire a known
directionality.

15 Without prejudice to the generality of the foregoing, the
invention includes, inter alia, the use of adaptored and
sorted sequences as PCR probes; the use of the present
method to facilitate investigation of a population of
sequences to identify a preselected fragment or sequence of
20 interest; the use of the present method to sort or
categorise (eg for sequencing purposes) sets of sequences
produced by exposing a nucleic acid source material to the
action of restriction endonucleases; the use of the present
method to increase the abundance of any given sequence in
25 a population thereby to facilitate investigation by a
hybridization probe; the use as a hybridization probe of a
sorted subset (or subsets) of sequences; the use of the

present method in clone identification or mapping; and, the use of the present method in examining in a species sequence polymorphisms, optionally monitored through more than one generation of the species in question.

5

It will be appreciated that the above is not exhaustive, and it should not be construed as limiting on the applicability of the present pioneering invention.

10 In other aspects, the invention obviously includes:

(a) The use of a population of adaptor molecules, each of such molecules carrying a nucleic acid sequence end recognition means, in categorizing or sorting a
15 nucleic acid sample into predetermined subsets of nucleic acid sequences, wherein each such adaptor molecule carries a nucleic acid sequence end recognition means which is specific to a predetermined base.

20

(b) A kit including reagents for use in a nucleic acid categorization process and comprising a population of adaptor molecules in the form of double stranded oligonucleotides having a single stranded end
25 extension as a nucleic acid sequence end recognition means.

- (c) A kit as above, wherein:
- (i) the adaptor molecule nucleic acid sequence end recognition means recognises a predetermined base, the presence of which base can form the basis for selection and/or
 - (ii) the primers (if present) include within their sequence a predetermined base, the presence of which base can form the basis for selection.
- (d) A population of adaptor molecules for use in a nucleic acid categorization process, wherein the adaptor molecules include nucleic acid sequence end recognition means recognising a predetermined base thereby permitting categorization of nucleic acid sequences linked to said adaptor molecules on the basis of selecting for a subset in which a chosen one of said predetermined bases has been recognised.
- (e) A process for the categorization of nucleic acid sequences in which said sequences are linked to a population of adaptor molecules each exhibiting specificity for linking to a sequence including a predetermined nucleotide base, categorization of the resulting linked sequences being based upon selection for the base.

Claims

1. A process for categorizing uncharacterized nucleic acid by sorting said nucleic acid into sequence-specific subsets, which process comprises:
- 5 (a) optionally, initially subjecting said uncharacterized nucleic acid to the action of a reagent, preferably an endonuclease, which reagent cleaves said nucleic acid so as to produce smaller size cleavage products;
- 10 (b) reacting either the uncharacterized nucleic acid or, as the case may be, cleavage products from (a) with a population of adaptor molecules so as to generate adapted products, each of which adaptor molecules carries nucleic acid sequence end recognition means,
- 15 and said population of adaptor molecules encompassing a range of such molecules having sequence end recognition means capable of linking to a predetermined subset of nucleic acid sequences; and
- 20 (c) selecting and separating only those adapted products resulting from (b) which include an adaptor of chosen nucleic acid sequence end recognition means.
2. A process as claimed in claim 1, wherein the adaptor molecules include oligonucleotides in which the nucleic acid sequence end recognition means comprises a single
- 25 stranded end of known nucleotide composition.

3. A process as claimed in claim 2, wherein the single stranded end exhibits complementarity to a predetermined nucleotide.

5 4. A process as claimed in any one of claims 1 to 3, wherein the population of adaptor molecules includes individual molecules such that both ends of the uncharacterised nucleic acid or cleavage products can be linked thereto.

10

5. A process as claimed in any one of claims 1 to 4, wherein at least some of the adaptor molecules are adapted to be immobilized on a solid phase.

15 6. A process as claimed in any one of claims 1 to 5, wherein selection of adapted products is based upon selecting those adaptor molecules having a single predetermined base.

20 7. A process as claimed in any one of claims 1 to 6, wherein optional step (a) is effected using an endonuclease specific to double stranded nucleic acid or having the capability of cutting at a recognised sequence on a single stranded nucleic acid.

25

8. A process as claimed in claim 7, wherein the endonuclease is selected from those which cleave single-

stranded nucleic acids, for example, BstNI, DdeI, HgaI, HinfI, or MnlI.

9. A process as claimed in any one of claims 1 to 8,
5 wherein at least some of the adaptor molecules also
comprise a known sequence permitting hybridization with a
PCR primer.

10. A process as claimed in claim 9, wherein selection of
10 adapted products is based upon selecting products
subjected to priming of nucleic acid synthesis in which a
primer has a single predetermined base, preferably at the
3' end.

15 11. A process of categorising uncharacterised nucleic
acid, which process comprises:

- (a) subjecting said uncharacterised nucleic acid to the
action of a reagent, preferably an endonuclease which
has cleavage and recognition site separated, which
20 reagent cleaves said nucleic acid so as to produce
double stranded cleavage products the individual
strands of which overlap at cleaved ends to leave a
single strand extending to a known extent;
- (b) ligating the cleavage products from (a) with adaptor
25 molecules to generate adapted cleavage products,
each of which adaptor molecules has a cleavage product
end recognition sequence and the thus-used adaptor

molecules encompassing a range of adaptor molecules having recognition sequences complementary to a predetermined subset of the sequences of the cleavage-generated extending single strands; and

- 5 (c) selecting and separating only those adapted cleavage products resulting from (b) which carry an adaptor of known recognition sequence.

12. A process as claimed in claim 11, wherein in step (b)
10 a number of separate reactions are performed in each of which a subset of the range of adaptor molecules is ligated.

13. A process as claimed in claim 11 or claim 12, wherein
15 the uncharacterized nucleic acid consists of single stranded nucleic acid which is first converted to double stranded nucleic acid.

14. A process as claimed in any one of claims 11 to 13,
20 wherein an endonuclease is employed in step (a) which is chosen from Class II restriction endonucleases the cleavage sites of which are asymmetrically spaced across the two strands of a double stranded substrate, and the specificity of which is not affected by the nature of the bases
25 adjacent to a cleavage site; optionally wherein the endonuclease is Fok 1.

15. A process as claimed in any one of claims 11 to 14, wherein the cleavage products from (a) have extending single strands which are from 1 to 10 bases in length, optionally from 4 to 6 bases.

5

16. A process as claimed in any one of claims 11 to 15, wherein the adaptor molecules have cleavage product end recognition sequences which comprise extending single strands which are complementary to the cleavage product
10 extending single strands resulting from step (a).

17. A process as claimed in claim 16, wherein at least some adaptor molecule cleavage product end recognition sequences end with a 5' hydroxyl group.

15

18. A process as claimed in any one of claims 11 to 17, wherein adaptor molecule cleavage product end recognition sequences are at least three nucleotides in length with unselected random bases present at each position two or
20 more nucleotides in from the end, optionally a 5' end.

19. A process as claimed in any one of claims 11 to 18, wherein the adaptor molecules also comprise a portion permitting separation when a nucleic acid sequence is
25 attached to its adaptor.

20. A process as claimed in claim 19, wherein the portion

permitting separation includes a biotin residue.

21. A process as claimed in any one of claims 10 to 20,
wherein selection of adapted products is based upon
5 selecting those adaptor molecules having a single
predetermined base.

22. A process as claimed in any one of claims 11 to 21,
wherein the adaptor molecules also comprise a predetermined
10 core sequence so that nucleic acid sequences which have
been linked to such molecules can be amplified by the use
of a primer exhibiting complementarity to the core
sequence.

23. A process as claimed in claim 22 wherein selection of
15 adapted products is based upon selecting products
subjected to priming of nucleic acid synthesis in which a
primer has a single predetermined base, preferably at the
3' end.

24. A process as claimed in any one of claims 11 to 23,
wherein adaptor molecules are employed in (b) which
20 correspond to all possible members of a chosen subset of
cleavage products present after step (a).

25. A process as claimed in any one of claims 11 to 24,
wherein at least some of the adaptor molecules carry a

predetermined recognition sequence which differs from that of all other adaptors present.

26. The use of a population of adaptor molecules, each of
5 such molecules carrying a nucleic acid sequence end
recognition means, in categorizing or sorting a nucleic
acid sample into predetermined subsets of nucleic acid
sequences, wherein each such adaptor molecule carries a
nucleic acid sequence end recognition means which is
10 specific to a predetermined base.

27. The use of claim 26, wherein the adaptor molecules are
double stranded oligonucleotides having a single stranded
extension portion at an end which serves as the nucleic
15 acid sequence end recognition means.

28. The use of claim 26 and 27, wherein the adaptor
molecules also include a sequence permitting hybridization
with a primer for nucleic acid amplification of any nucleic
20 acid sequence linked to the adaptor molecules.

29. A kit including reagents for use in a nucleic acid
categorization process and comprising a population of
adaptor molecules in the form of double stranded
25 oligonucleotides having a single stranded end extension as
a nucleic acid sequence end recognition means.

30. A kit as claimed in claim 29 also including primers for use in a nucleic acid amplification process, at least some of the adaptor molecules including a sequence for primer hybridization thereto.

5

31. A kit as claimed in claim 29 or claim 30, wherein:

- (a) the adaptor molecule nucleic acid sequence end recognition means recognises a predetermined base, the presence of which base can form the basis for selection and/or
- (b) the primers (if present) include within their sequence a predetermined base, the presence of which base can form the basis for selection.

10

32. A kit as claimed in any one of claims 29 to 31, wherein the adaptor molecules include nucleotide sequence which, in use, not only permits indexing of nucleic acid sequences linked to said adaptor molecules but also establishes directionality in said nucleic acid sequences.

20

33. A kit as claimed in any one of claims 29 to 32, wherein at least some of the adaptor molecules also include means permitting immobilization on a solid phase of such adaptor molecules when linked to nucleic acid sequences.

25

34. A kit as claimed in any one of claims 29 to 33, and also including a nucleic acid cleavage reagent, optionally

an endonuclease, for example Fok 1.

35. A population of adaptor molecules for use in a nucleic acid categorization process, wherein the adaptor molecules include nucleic acid sequence end recognition means recognising a predetermined base thereby permitting categorization of nucleic acid sequences linked to said adaptor molecules on the basis of selecting for a subset in which a chosen one of said predetermined bases has been recognised.

36. A process for the categorization of nucleic acid sequences in which said sequences are linked to a population of adaptor molecules each exhibiting specificity for linking to a sequence including a predetermined nucleotide base, categorization of the resulting linked sequences being based upon selection for the base.

37. The use of the steps of a process as claimed in any one of claims 1, 11 or 36 in any one of the following:

- (a) enhancing the amount of a chosen nucleic acid sequence from a mixture of nucleic acid sequences;
- (b) investigating polymorphisms in nucleic acid sequences;
- (c) sequencing a nucleic acid sequence;
- (d) facilitating the use of nucleic acid hybridization probes by enhancing the amount of selected sequences present in a sample being probed;

- 5 (e) increasing the number of different nucleic acid sequences which can be used as a probe by employing separately as probes subsets of the original multisequence probe produced by categorization thereof;
- (f) fingerprinting the individual members of nucleic acid libraries for clone identification or mapping;
- (g) producing a set of adapted and sorted sequences for use as PCR probes;
- 10 (h) identification within a population of nucleic acid sequences of a preselected fragment or sequence of interest;
- (i) comparing nucleic acid sequence frequency in two or more nucleic acid sequence population samples.



Figure 1.**Basic scheme for selecting specific restriction fragments.**

Figure 2.

Nature of the cleavage produced by the restriction endonuclease Fok 1.



n - can be any base but is always the same base at any point in a given sequence of DNA.

↓ sites of cleavage

Figure 3a

Base specific adaptor at a Fok 1 cleavage site - specificity introduced at the two overhanging 5' bases of the adaptor




Figure 3b

Selection for fragments produced by type II restriction endonucleases whose cuts are staggered and do not overlap the recognition site.

1. Base specific adaptor and isolation of adapted fragments.




Figure 4.

Examples of adapters and primers used in the sorting process.Biotinylated universal adaptor for specific adaptors followed by specific adaptor options

5' Bio GTTCTCGGAGCACTGTCCGAGA 3'
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄AA 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄AC 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄AG 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄AT 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄CA 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄CC 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄CG 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄CT 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄GA 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄GC 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄GG 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄GT 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄TA 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄TC 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄TG 5'
 or
 3' CAAGAGCCTCGTGACAGGCTCT(N)₄(N)₄TT 5'

Non specific adaptor combination

3'(N)₄(N)₄(N)₄(N)₄AGGAAGAGGACGCTGTCTGT 5'
 5' TCCTTCTCCTGCGACAGACA 3'

Possible primers for use with non specific adaptor combination

3'AAGGAAGAGGACGCTGTCTGT 5'
 or
 3'CAGGAAGAGGACGCTGTCTGT 5'
 or
 3'GAGGAAGAGGACGCTGTCTGT 5'
 or
 3'TAGGAAGAGGACGCTGTCTGT 5'

Primer options for use with specific adaptors

5' GTTCTCGGAGCACTGTCCGAGAA 3'
 or
 5' GTTCTCGGAGCACTGTCCGAGAC 3'
 or
 5' GTTCTCGGAGCACTGTCCGAGAG 3'
 or
 5' GTTCTCGGAGCACTGTCCGAGAI 3'

6/8



Figure 5a**Base specific priming of base specifically adapted Fok 1 fragment**


Figure 5b

Selection for fragments produced by type II restriction endonucleases whose cuts are staggered and do not overlap the recognition site.
2. Base specific priming of DNA synthesis.



8/8

Figure 6



INTERNATIONAL SEARCH REPORT

International Application No.

PCT/GB 93/01452

I. CLASSIFICATION OF SUBJECT MATTER (In several classification symbols apply, indicate all) * According to International Patent Classification (IPC) or to both National Classification and IPC IPC ⁵ : C 12 Q 1/68		
II. FIELDS SEARCHED		
Minimum Documentation Searched †		
Classification System	Classification Symbols	
IPC ⁵	C 12 Q	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ‡		
Category *	Citation of Document, †† with indication, where appropriate, of the relevant passages ††	Relevant to Claim No. ††
A	EP, A2, 0 405 376 (BOEHRINGER INGELHEIM INTER- NATIONAL GMBH) 02 January 1991 (02.01.91), claim 1.	1, 7-9, 11, 14, 37
A	Chemical Abstracts, vol. 109, no. 11, issued 1988, September 12 (Columbus, Ohio, U.S.A.), S.C. KIM et al. "Cleaving DNA at any predetermined site with adapter-primers and class-IIS restriction enzymes.", page 197, the abstract no. 87440j, Science (Washington, D.C., 1988, 240(4851), 504-6.	1, 8, 14
A	Chemical Abstracts, vol. 109, no. 20, issued 1988,	1, 8, 14
* Special categories of cited documents: †† "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed † later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docu- ments, such combination being obvious to a person skilled in the art. "Z" document member of the same patent family		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
22 September 1993		03.11.93
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		SCHNASS e.h.

Form PCT/ISA/710 (second sheet) (January 1985)

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, " with indication, where appropriate, of the relevant passages	Relevant to Claim No.
A	<p>November 14 (Columbus, Ohio, U.S.A.), W. MANDECKI et al. "FokI method of gene synthesis.", page 210, the abstract no. 184916b, Gene 1988, 68(1), 101-7.</p> <p>---</p>	1,5
	<p>Chemical Abstracts, vol. 115, no. 7, issued 1991, August 19 (Columbus, Ohio, U.S.A.), L. HALL et al. "A rapid and efficient method for site-directed mutagenesis by PCR, using biotinylated universal primers and streptavidin-coated magnetic beads.", page 253, the abstract no. 66165g, Protein. Eng. 1991, 4(5), 601.</p> <p>----</p>	

ANHANG

zum internationalen Recherchen-
bericht über die internationale
Patentanmeldung Nr.

ANNEX

to the International Search
Report to the International Patent
Application No.

ANNEXE

au rapport de recherche inter-
national relatif à la demande de brevet
international n°

PCT/GB 93/01452 SAE 76533

In diesem Anhang sind die Mitglieder
der Patentfamilien der in obenge-
nannten internationalen Recherchenbericht
angeführten Patentdokumente angegeben.
Diese Angaben dienen nur zur Unter-
richtung und erfolgen ohne Gewähr.

This Annex lists the patent family
members relating to the patent documents
cited in the above-mentioned inter-
national search report. The Office is
in no way liable for these particulars
which are given merely for the purpose
of information.

La présente annexe indique les
membres de la famille de brevets
relatifs aux documents de brevets cités
dans le rapport de recherche inter-
national visée ci-dessus. Les renseigne-
ments fournis sont donnés à titre indica-
tif et n'engagent pas la responsabilité
de l'Office.

In Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
EP A2 405376	02-01-91	CA AA 2019663 EP A3 405376 FI A0 903143 JP A2 3117499 DE A1 4020028	24-12-90 25-11-92 21-06-90 20-05-91 03-01-91

